

Remarkable Mass Relations in the Electroweak Model

Jean Pestieau

Institut de Physique Théorique, Université catholique de Louvain,
Chemin du Cyclotron 2, B-1348 Louvain-la-Neuve, Belgium¹

Abstract

In the electroweak standard model we observe two remarkable empirical mass relations, $m_W + m_B = \frac{v}{2}$ and $m_W - m_B = e\frac{v}{2}$ where $m_Z^2 \equiv m_W^2 + m_B^2$, e is the positron electric charge and v , the strength of the Higgs condensate.

¹pestieau@fyma.ucl.ac.be

The electroweak standard model [1, 2, 3] is based on the gauge group $SU(2) \otimes U(1)$, with gauge bosons fields $W_\mu^i = 1, 2, 3$ and B_μ . After spontaneous symmetry breaking, W_μ^i and B_μ acquire masses, m_W and m_B respectively. W_μ^3 and B_μ are mixed. After diagonalization of mass matrix, one gets the physical fields A_μ and Z_μ corresponding to the massless photon and the neutral Z boson of mass m_Z , with the relations [2, 4]

$$m_Z^2 = m_W^2 + m_B^2 \quad (1)$$

$$\cos \theta_W = \frac{m_W}{m_Z}, \quad \sin \theta_W = \frac{m_B}{m_Z} \quad (2)$$

where θ_W is the weak mixing angle :

$$Z_\mu = W_\mu^3 \cos \theta_W - B_\mu \sin \theta_W \quad (3)$$

$$A_\mu = W_\mu^3 \sin \theta_W + B_\mu \cos \theta_W \quad (4)$$

From experimental data, we get two remarkable empirical mass relations defining two different mass scales :

$$m_W + m_B = \frac{v}{2} \quad (5)$$

$$m_W - m_B = e \frac{v}{2} \quad (6)$$

where e is the positron electric charge and v , the strength of the Higgs condensate.

Eqs (2), (5) and (6) imply [5] :

$$e = \frac{1 - \operatorname{tg} \theta_W}{1 + \operatorname{tg} \theta_W} = \operatorname{tg} \left(\frac{\pi}{4} - \theta_W \right) \quad (7)$$

From the experimental data [6]

$$\alpha = \frac{e^2}{4\pi} = \frac{1}{137.03599976(50)} \quad (8)$$

$$m_Z = 91.1872 \pm 0.0021 \text{ GeV}, \quad (9)$$

we obtain :

$$\operatorname{tg} \theta_W = 0.5351 \quad (10)$$

$$m_W = \frac{v}{4}(1 + e) = 80.400 \pm 0.002 \text{ GeV} \quad (11)$$

$$m_B = \frac{v}{4}(1 - e) = 43.022 \pm 0.001 \text{ GeV} \quad (12)$$

$$v = 246.85 \text{ GeV} \quad (13)$$

to be compared with the experimental values [6] :

$$m_W(\text{exp}) = 80.419 \pm 0.038 \text{ GeV} \quad (14)$$

$$v_F \equiv \left(\frac{1}{\sqrt{2}G_F} \right)^{1/2} = 246.22 \text{ GeV} \quad (15)$$

v_F is the usual value of the strength of the Higgs condensate obtained from G_F , the Fermi constant derived from the muon lifetime [6] and

$$\frac{v}{v_F} - 1 = 0.0026 \quad (16)$$

We conclude that Eqs (5) and (6) are robust relations and point to new physics.

With respect to weak $SU(2)$, W_μ^i 's transform as a weak isotriplet, B_μ as a weak singlet. Let us define

$$D_\mu = \frac{1}{\sqrt{2}}(W_\mu^3 - B_\mu) \quad (17)$$

$$U_\mu = \frac{1}{\sqrt{2}}(W_\mu^3 + B_\mu). \quad (18)$$

W_μ^+ , W_μ^- , D_μ and U_μ are the fields obtained from the direct product of two isodoublet fundamental fields. Then the masses of D and U fields are

$$m_D = \frac{1}{\sqrt{2}}(m_W + m_B) = \frac{v}{2\sqrt{2}} = 87.27 \text{ GeV} \quad (19)$$

$$m_U = \frac{1}{\sqrt{2}}(m_W - m_B) = \frac{ev}{2\sqrt{2}} = 26.43 \text{ GeV}. \quad (20)$$

Now we can see the pattern of the neutral vector boson mass matrix :

$$\begin{aligned} 2\mathcal{L}_M &= \widehat{Z_\mu A_\mu} \begin{pmatrix} m_Z^2 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} Z_\mu \\ A_\mu \end{pmatrix} = \\ &\widehat{W_\mu^3 B_\mu} \begin{pmatrix} m_W^2 & -m_W m_B \\ -m_W m_B & m_B^2 \end{pmatrix} \begin{pmatrix} W_\mu^3 \\ B_\mu \end{pmatrix} = \end{aligned}$$

$$\begin{aligned}
\widehat{D_\mu U_\mu} &= \begin{pmatrix} m_D^2 & m_D m_U \\ m_D m_U & m_U^2 \end{pmatrix} \begin{pmatrix} D_\mu \\ U_\mu \end{pmatrix} \\
\widehat{D_\mu U_\mu} &= \frac{v^2}{8} \begin{pmatrix} 1 & e \\ e & e^2 \end{pmatrix} \begin{pmatrix} D_\mu \\ U_\mu \end{pmatrix}
\end{aligned} \tag{21}$$

It is an empirical fact that the electric charge sets the mass scale between U_μ and D_μ fields:

$$e = \frac{m_U}{m_D} = \operatorname{tg} \left(\frac{\pi}{4} - \theta_W \right). \tag{22}$$

It is interesting to note that

$$m_t = 174.3 \pm 5.1 \text{ GeV} \tag{23}$$

to be compared with the empirical relation

$$m_t = \frac{v}{\sqrt{2}} = \sqrt{2}(m_W + m_B) = 174.6 \text{ GeV} \tag{24}$$

and comparing Eqs (24) and (19), we find

$$\frac{m_t}{m_D} = 2 \tag{25}$$

References

- [1] S.L. Glashow, *Nucl. Phys.* **22** (1961) 579.
- [2] S. Weinberg, *Phys. Rev. Lett.* **19** (1967) 1264.
- [3] A. Salam, in *Elementary Particle Physics: Relativistic Groups and Analyticity*, Ed. N. Svartholm (Almqvist and Wiksell, 1968).
- [4] J. Pestieau and P. Roy, *Phys. Rev. Lett.* **23** (1969) 349; *Lett. Nuovo Cimento* **31** (1981) 625.
- [5] G. Lopez-Castro and J. Pestieau, hep-ph/9804272.
- [6] *Review of Particle Physics*, *Eur. Phys. J.* **C15** (2000) 1.